

RESEARCH ARTICLE

The Levels of Vitamin E in Seminal Fluids and their Association with Male Infertility: Short Review

Abdulsamie H. Alta'ee^{1,2}, Hanan Abbas Majeed³, Mahmoud H. Hadwan^{4*}

¹College of Medicine, University of Babylon, Postal Code 51002, Hilla P.O. Box 473, Iraq.

²College of Pharmacy, University of Babylon, Postal Code 51002, Hilla P.O. Box 4, Iraq.

³Community Health Department, Alfurat Alawsat Technical University, Technical Institute of Karbala, Karbala City; Iraq.

⁴Chemistry Department, College of Science, University of Babylon, Postal Code 51002, Hilla P.O. Box 4, Iraq.

*Corresponding Author E-mail: mahmouhadwan@gmail.com

ABSTRACT:

Background: Vitamin E is one of four fat-soluble vitamin with antioxidant property consists of tocopherols and tocotrienols with α -tocopherol the most bioactive member of this family. Aim: This study aims to assess always almost each potential deals with the effect of consuming vitamin E on seminal fluids properties in asthenospermic patients. **Methods:** This review reports the modern state of knowledge and ambiguities about the oral treatment of infertile male with vitamin E. The current study illustrates the best obtainable information from randomized clinical trials, investigational studies, and systematic reviews. **Results:** Data extracted from the previous publications clarify that treatment of idiopathic infertile males with vitamin E supplementation can lead to a considerable enhancement in semen quality; however large-scale studies will be of great benefit to achieve such conclusions. **Conclusions:** The vitamin E therapy is an effective choice for the management of idiopathic infertile males with asthenozoospermia.

KEYWORDS: Vitamin E, infertility, asthenozoospermia, oxidative stress; antioxidant.

INTRODUCTION:

World Health Organization (WHO) describes infertility as the "disease of the reproductive system defined by the failure to achieve a clinical pregnancy after 12 months or more of regular unprotected sexual intercourse¹. Infertility affects approximately 15% couples in the reproductive-aged worldwide, half of this percent attributed to infertility male factors, of which there are about 50% of cases yet of idiopathic origin^{2,3}.

Oxidative stress and infertility:

Oxidative stress (OS) is a critical pathophysiological factor in the causation of a male infertility of idiopathic origin⁴. It is a consequence of increase generation of reactive oxygen and/or nitrogen species which collectively called free radicals, or produced by decreased antioxidant production. The balance between reactive oxygen species (ROS) and/or reactive nitrogen species (RNS) with antioxidants is strictly maintained in a fertile male, and their imbalance is implicated in various pathological conditions concern sperm morphology and function, as well as their adverse effect on sperm genomic integrity⁵. OS is highly susceptible to induce damage in spermatozoa due to the presence of high content of polyunsaturated fatty acids (PUFA) in the seminal plasma membrane, low levels of seminal defense enzymes⁶ and high levels of free radicals⁷. OS disables normal sperm function and induces mitochondrial genome mutations by the action of peroxidation process to PUFA especially docosahexaenoic acid; one of the components of sperm plasma membrane^{8,9}. The lipid peroxides formation

affects sperm membrane fluidity and flagellar motion, leading to sperm dysfunction. Mitochondrial mutations impair metabolism within spermatozoa resulting in an energy crisis, which in turn lead to the spermatozoa functional impairment¹⁰.

Progress of infertility:

Accumulating medical reports confirmed a great decrease in sperm count in the last fifty years, where Carlsenet *al.*,¹¹ found after the analysis of data of the mean seminal volume and sperm density in 14947 males cases subjected to sixty-one studies extended from 1938 to 1991. Carlsenet *al.*,¹¹ study found a significant depletion in seminal quantity from 3.4 ml to 2.75 ml and sperm density from 113 million/ml to 66 million/ml in the comparison between 1940 and 1990. This shows that there is 20% decrease in seminal volume and 58% decrease in sperm density in the last five decades. Other studies confirmed this results carried out in the last two decades also show a decrease in semen quality and they attributed their finding to the multifactor such as diet, lifestyle and environmental factors which may in combination interfere with spermatogenesis^{12,13}.

Chemistry of vitamin E:

Fat soluble vitamins like vitamin E and A which are naturally occurring in diet have a preservative properties make them ideal excipients materials used many industries including in drug industries¹⁴. The name vitamin E is colligative term involves eight relative compounds, each of them consists of a phytol side chain attach to chromanol ring; four tocotrienols compounds categorized as (α , β , γ , and δ) and four tocopherols compounds categorized as (α , β , γ , and δ)¹⁵. The prefixes α , β , γ , and δ attach to tocopherols give to distinguish among the position of methyl groups substituted on chromanol ring¹⁶. α -Tocopherol is the widely distributed among natural sources and is the most active member of tocopherol family and has the ability to quench a couple of peroxy radicals that responsible for triggering lipid peroxidation process¹⁷. For this reason vitamin E protect lipid found in membrane against free radical attack¹⁸, and thus stabilize membrane¹⁹. The digestion of vitamin E is analogous to that of carotenoids and vitamin A, and the deficiency in its level may occur in fat malabsorption case with neurological symptoms. These symptoms result from the decrease in nerve conduction, which may be reversed by E vitamin supplementation¹⁷.

The absorption of vitamin E occurs in the gut via lymph system and intestine via chylomicron to reach liver as chylomicron remnants. It is then distributed by very low density lipoprotein (VLDL) to required tissue, and can

transfer to low density lipoprotein (LDL) or scavenges by high density lipoprotein (HDL)²⁰.

Among deferent types of tocopherol, α -tocopherol represents the predominant and more active form of vitamin E that present in the leaves of plants with an antioxidant property. Several studies were reported that vitamin E plays a crucial role in the protection of plant photosynthetic apparatus against oxidative stress^{21,22}. Vitamin E also has a key role in the amino acids and lipids transportation in the intestine²². Other function of vitamin E includes its role in steroidogenesis and iron metabolism²³, as well as its role in the stimulation of hormonal response and the response of cellular immune system against infection threatening²⁴. Vitamin E deficiency symptoms and disorders are vary depending on the species affected^{25,26}.

Vitamin E as treatment:

Vitamin E represents the first line of the chain-breaking antioxidant present in the cell membrane. It is well-documented as a lipid-soluble antioxidant and it has an inhibitory effect on ROS and/or RNS that may produce damage when attack cell membrane including seminal membrane. The product of lipid peroxidation process as result of oxidative stress in the seminal specimens is malondialdehyde (MDA) found to be significantly decreased after vitamin E supplementation, as well as notable enhancement in sperm motility²⁷. A clinical trial carried out by Kessopoulouet *al.*²⁸ shows that supplementation of 600 mg per day of vitamin E enhances sperm function, in the term of the sperm ability to penetrate the female egg in vitro. Another clinical trial was reported that after taken vitamin E in combination with the trace element selenium, the volunteers status with altered sperm motility and low sperm count were improved after four months of an experiment in the term of total percent of normal spermatozoa and sperm motility²⁹.

The role of vitamin E in infertility:

Vitamin E has an advantage of competing for peroxy radicals much faster than PUFA, therefore, a small amount of vitamin E is sufficient to protect a large amount of PUFA against oxidative stress. It has been found that one molecule of vitamin E corresponds to thousands of lipid molecules in biological membranes³⁰. During this process, vitamin E transformed into vitamin E radical which unless is regenerated by other chain breaker antioxidant like vitamin C, it must be replenished from diet or from body storage, or it will be consumed eventually³¹, as shown in Figure 1.




Figure 1 Mechanism of antioxidant defense system. Modified from (Alta'ee et al., 2015) (R= free radical; PUFA-OO=peroxy radical of PUFA; PUFA-OOH=hydroperoxy PUFA; PUFA-OH=hydroxyPUFA; vitamin E'= vitamin E radical; GSH= reduced glutathione; GS-SG= oxidized glutathione; Se= selenium).

Vitamin E as treatment:

Many organs and tissues like adipose tissue may reveal a deficiency of vitamin E accompanied with a number of disorders such as ischemic heart disease, male infertility, breast cancer and increase the predisposing to infections^{16,32}, as well as the promotion of environmental and dietary stress in both animals and humans^{22,24,33}. Animal clinical trial show low toxicity of vitamin E and no carcinogenic, mutagenic, or teratogenic effect in comparison to human clinical trials which were found many symptoms associated with vitamin E toxicity such as nausea, stupor, and loss of appetite, as well as enhancement of calcium absorption, which in turn may lead to increase deposition and precipitation of Ca in several organs like kidney and arteries²¹.

Studies carried out on human and animals show a positive effect for vitamin E on fertility as a dietary essential factor in both genders^{33,34}. In spite of the fact that vitamin E has multifunction in the body, it possesses a characteristic role in the fertility of human and many types of animals species. Poultry farmers add vitamin E to the feed to enhance male and female fertility, which in turn increases the net income of poultry production³⁵.

Numerous studies reported the antioxidant properties of vitamin E for improvement of sperm motility and count^{36,37,38}. Other studied demonstrated positive effect for the usage of high doses of vitamin E on improving fertility^{28,29}, whereas opposite result is obtained by another study, in which no such advantage for the usage of high doses of combination of vitamin C and E³⁹.

Male with idiopathic infertility treated with vitamin E found to have a rate of 15% of seminal live birth compared with 0% of the control group in meta-analysis

clinical trials^{28,40,41}. However, the usage of different doses of vitamin E in these studies range from 300 mg to 600 mg making the determination of optimum dose is a matter under discussion. Moreover, interesting results were presented by Rolf et al.,⁴¹ study which investigates the defense mechanism of vitamin E as an antioxidant in infertile men after treating with a high dose of E vitamin (800 mg) for a period of eight weeks. This randomized, placebo-controlled trial advice to measure the seminal ROS levels before and after treating with vitamin E, due to the uselessness of this type of therapy in infertile men don't suffer from oxidative stress. This is very important because the high level of vitamin E may act as pro-oxidant in the semen of infertile men without oxidative stress⁴².

Several reports focused upon the combinations of the mixture of antioxidants in the treatment of idiopathic infertile males. The pharmaceutical companies also started to produce combinations of such antioxidants as the mixture of vitamin C and vitamin E in the therapy of such patients. However, the data of such reports found to be fluctuated, where⁴³ study which uses a daily pill of one gram of vitamin E and one gram vitamin C for two months for treatment of infertile males was found a significant reduction in DNA fragmentation in semen of infertile males when compared with placebo group, however, this study found no effect on other seminal fluid parameters. Another study⁴¹ was found a comparable result, in which one gram of vitamin C and 800 gram of vitamin E were used daily for eight weeks. This study didn't find an improvement in seminal fluid parameters in males and the rate of pregnancy in females. Another study⁴⁴ use the same combination in addition to vitamin A and found the same results in the idiopathic infertile men. In a placebo-control clinical

trial carried out on sixty idiopathic infertile men, the combination of 400 mg vitamin E and 25 mg clomiphene citrate was used daily for six months, and the results shown significant improvement in sperm count and motility in treated group in the comparison group⁴⁵. Patankar et al.⁴⁶ investigates the role of clomiphene citrate on primary infertile males and shows significant improvement in seminal fluid parameters. Furthermore, the supplemented of E vitamin was studied alone in another clinical trial and this study was also shows a significant improvement in seminal fluid analysis and the pregnancy rate²⁷. The usage of combination therapy of vitamin E with other antioxidants were studied in six clinical trials^{41,44,47-50} and show significant improvement of sperm motility in the first three studies and significant improvement of sperm count in one of them^{50,51}.

From all of above, the subject of vitamin E supplementation for the treatment of infertility of both genders is a controversial issue that needs further investigation in studies that extend for longer time and require a larger sample size.

CONCLUSIONS:

The present literature review shows that E vitamin is required for the function and development of the reproductive tissues in both sexes, possibly due to its key role in the modulation of antioxidant balance. The excessive generation of ROS makes biological systems subject to continuous exposure to oxidative stress. The excessive production ROS may contribute in several pathogenic conditions including cancer, diabetes mellitus, aging, and several diseases affecting reproduction systems in both sexes. The presence of active antioxidant defense mechanism may contribute to prevent or decrease the oxidative induced injury.

REFERENCES:

1. Patricio A, Cruz DF, Silva JV, Padrão A, Correia BR, Korrodi-Gregório L, Ferreira R, Maia N, Almeida S, Lourenço J, Silva V. Relation between seminal quality and oxidative balance in sperm cells. *Acta Urológica Portuguesa*. 2016 Apr 30;33(1):6-15.
2. Olayemi FO. review on some causes of male infertility. *African Journal of Biotechnology*. 2010;9(20).
3. Wong, Wai Yee, et al. "Male factor subfertility: possible causes and the impact of nutritional factors." *Fertility and sterility* 73.3 (2000): 435-442.
4. Köksal IT, Tefekli A, Usta M, Erol H, Abbasoglu S, Kadioglu A. The role of reactive oxygen species in testicular dysfunction associated with varicocele. *BJU international*. 2000 Sep 1;86(4):549-52.
5. Shiraishi K, Naito K. Generation of 4-hydroxy-2-nonenal modified proteins in testes predicts improvement in spermatogenesis after varicocelectomy. *Fertility and sterility*. 2006 Jul 31;86(1):233-5.
6. Alsalmán AR, Almashhedy LA, Hadwan MH. Zinc supplementation attenuates lipid peroxidation and increases antiperoxidant activity in seminal plasma of Iraqi asthenospermic men. *Life Sci J*. 2013;10(4):989-97.
7. Giacone F, Condorelli RA, Mongioì LM, Bullara V, La Vignera S, Calogero AE. In vitro effects of zinc, D-aspartic acid, and coenzyme-Q10 on sperm function. *Endocrine*. 2017 May 1;56(2):408-15.
8. Hadwan MH, Almashhedy LA, Alsalmán AR. Oral zinc supplementation restores superoxide radical scavengers to normal levels in spermatozoa of Iraqi asthenospermic patients. *International Journal for Vitamin and Nutrition Research*. 2015 Jan 1;85(3-4):165-73.
9. Hadwan MH, Almashhedy LA, Alsalmán AR. Seminal xanthine oxidase: Appropriate fluorometric assay for the examination of spermatozoa disorders. *BioTechnology: An Indian Journal*. 2014;9(7).
10. Atig F, Kerkeni A, Saad A, Ajina M. Effects of reduced seminal enzymatic antioxidants on sperm DNA fragmentation and semen quality of Tunisian infertile men. *Journal of assisted reproduction and genetics*. 2017 Mar 1;34(3):373-81.
11. Carlsen E, Giwercman A, Keiding N, Skakkebaek NE. Evidence for decreasing quality of semen during past 50 years. *Bmj*. 1992 Sep 12;305(6854):609-13.
12. Van Waelegheem K, De Clercq N, Vermeulen L, Schoonjans FR, Comhaire F. Deterioration of sperm quality in young healthy Belgian men. *Human Reproduction*. 1996 Feb 1;11(2):325-9.
13. Sinclair S. Male infertility: nutritional and environmental considerations. *Alternative medicine review: a journal of clinical therapeutic*. 2000 Feb;5(1):28-38.
14. Jones AM, Baker R, Ragone D, Murch SJ. Identification of pro-vitamin A carotenoid-rich cultivars of breadfruit (*Artocarpus, Moraceae*). *Journal of food composition and analysis*. 2013 Aug 31;31(1):51-61.
15. Górnas P. Unique variability of tocopherol composition in various seed oils recovered from by-products of apple industry: Rapid and simple determination of all four homologues (α , β , γ and δ) by RP-HPLC/FLD. *Food chemistry*. 2015 Apr 1;172:129-34.
16. Hinchá DK. Effects of α -tocopherol (vitamin E) on the stability and lipid dynamics of model membranes mimicking the lipid composition of plant chloroplast membranes. *FEBS letters*. 2008 Oct 29;582(25-26):3687-92.
17. Brigelius-Flohe R, Traber MG. Vitamin E: function and metabolism. *The FASEB Journal*. 1999 Jul 1;13(10):1145-55.
18. Niki E, Yamamoto Y, Komuro E, Sato K. Membrane damage due to lipid oxidation. *The American journal of clinical nutrition*. 1991 Jan 1;53(1):201S-5S.
19. Srivastava S, Phadke RS, Govil G, Rao CN. Fluidity, permeability and antioxidant behaviour of model membranes incorporated with α -tocopherol and vitamin E acetate. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 1983 Oct 12;734(2):353-62.
20. Burton GW, Traber MG. Vitamin E: antioxidant activity, biokinetics, and bioavailability. *Annual review of nutrition*. 1990 Jul;10(1):357-82.
21. Munne-Bosch S. The role of α -tocopherol in plant stress tolerance. *Journal of plant physiology*. 2005 Jul 1;162(7):743-8.
22. Wang X, Quinn PJ. The location and function of vitamin E in membranes. *Molecular membrane biology*. 2000 Jan 1;17(3):143-56.
23. MacDonald RC, MacDonald RI, Menco BP, Takeshita K, Subbarao NK, Hu LR. Small-volume extrusion apparatus for preparation of large, unilamellar vesicles. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 1991 Jan 30;1061(2):297-303.
24. Oliver AE, Hinchá DK, Crowe LM, Crowe JH. Interactions of arbutin with dry and hydrated bilayers. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 1998 Mar 6;1370(1):87-97.
25. Baldi S, Innocenti M, Frascerra S, Nannipieri M, Lippi A, Rindi P, Ferrannini E. Effects of hemodialysis and vitamin E supplementation on low-density lipoprotein oxidizability in end-stage renal failure. *Journal of nephrology*. 2012 Dec;26(3):549-55.

26. Habibian M, Ghazi S, Moeini MM, Abdolmohammadi A. Effects of dietary selenium and vitamin E on immune response and biological blood parameters of broilers reared under thermoneutral or heat stress conditions. *International journal of biometeorology*. 2014 Jul 1;58(5):741-52.
27. Suleiman SA, Ali ME, Zaki ZM, El-Malik EM, Nasr MA. Lipid peroxidation and human sperm motility: protective role of vitamin E. *Journal of andrology*. 1996 Sep 10;17(5):530-7.
28. Kessopoulou E, Powers HJ, Sharma KK, et al. A double-blind randomized placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility. *FertilSteril* 1995;Oct64(4):825-831
29. Vézina D, Mauffette F, Roberts KD, Bleau G. Selenium-vitamin E supplementation in infertile men. Effects on semen parameters and micronutrient levels and distribution. *Biological Trace Element Research*. 1996 Jun 1;53(1):65-83.
30. Burton GW, Joyce A, Ingold KU. Is vitamin E the only lipid-soluble, chain-breaking antioxidant in human blood plasma and erythrocyte membranes?. *Archives of Biochemistry and Biophysics*. 1983 Feb 15;221(1):281-90.
31. Zennaro L, Rossetto M, Vanzani P, De Marco V, Scarpa M, Battistin L, Rigo A. A method to evaluate capacity and efficiency of water soluble antioxidants as peroxy radical scavengers. *Archives of biochemistry and biophysics*. 2007 Jun 1;462(1):38-46.
32. Matringe M, Ksas B, Rey P, Havaux M. Tocotrienols, the unsaturated forms of vitamin E, can function as antioxidants and lipid protectors in tobacco leaves. *Plant physiology*. 2008 Jun 1;147(2):764-78.
33. Khan RU, Naz S, Nikousefat Z, Tufarelli V, Javdani M, Rana N, Laudadio V. Effect of vitamin E in heat-stressed poultry. *World's poultry science journal*. 2011 Sep;67(3):469-78.
34. Khan RU, Rahman ZU, Nikousefat Z, Javdani M, Laudadio V, Tufarelli V. Vitamin E: pharmaceutical role in poultry male fecundity. *World's Poultry Science Journal*. 2012 Mar;68(1):63-70.
35. Ordas B, Vahedi S, Seidavi A, Rahati M, Laudadio V, Tufarelli V. Effect of Testosterone Administration and Spiking on Reproductive Success of Broiler Breeder Flocks. *Reproduction in domestic animals*. 2015 Oct 1;50(5):820-5.
36. Madding CI, Jacob M, Ramsay VP, Sokol RZ. Serum and semen zinc levels in normozoospermic and oligozoospermic men. *Ann NutrMetab* 1986;30:213-218.
37. Netter A, Hartoma R, Nahoul K. Effect of zinc administration on plasma testosterone, dihydrotestosterone, and sperm count. *Arch Androl* 1981;7:69-73.
38. Tikkiwal M, Ajmera RL, Mathur NK. Effect of zinc administration on seminal zinc and fertility of oligospermic males. *Indian J PhysiolPharmacol* 1987;31:30-34.
39. Lenzi A, Lombardo F, Gandini L, et al. Glutathione therapy for male infertility. *Arch Androl* 1992;29:65-68.
40. Showell MG, Brown J, Yazdani A, et al. Antioxidants for male subfertility. *Cochrane Database Syst Rev* 2011(1):CD007411
41. Rolf A, Cooper TG, Yeong CH, Nieschlag E. (1999) Antioxidants treatment of patients with asthenozoospermia or moderate oligoasthenozoospermia with high dose vitamin C and vitamin E. *Hum Reprod* 14, 1028–1033.
42. Hamada AJ, Montgomery B, Agarwal A. Male infertility: a critical review of pharmacologic management. *Expert opinion on pharmacotherapy*. 2012 Dec 1;13(17):2511-31.
43. Greco E, Iacobelli M, Rienzi L, et al. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl* 2005;May-Jun26(3):349-53
44. Scott R, MacPherson A & Yates RW. (1998) The effect of oral selenium supplementation on human sperm motility. *Br J Urol* 82, 76–80.
45. Ghanem H, Shamloul R. An evidence-based perspective to the medical treatment of male infertility: a short review. *Urologiainternationalis*. 2009;82(2):125-9.
46. Patankar SS, Kaore SB, Sawaneh MV, Mishra NV, Deshkari AM. Short communication effect of clomiphene citrate on sperm density in male partners of infertile couples. *Indian J PhysiolPharmacol*. 2007;51(2):195-8.
47. Keskes-Ammar L, Feki-Chakroun N, Rebai T, Sahnoun Z, Ghazzi H, Hammami S, Zghal K, Fki H, Damak J, Bahloul A. (2003) Sperm oxidative stress and the effect of an oral vitamin E and selenium supplement on semen quality in infertile men. *Arch Androl* 49, 83–94.
48. Omu AE, Al-Azemi MK, Kehinde EO, Anim JT, Oriowo MA & Mathew TC. (2008) Indication of the mechanisms involved in improved sperm parameter by zinc therapy. *Med PrincPract* 17, 108–116.
49. Greco E, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Tesarik J. (2005) Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl* 26, 349–353.
50. Galatioto GP, Gravina GL, Angelozzi G, Sacchetti A, Innominato PF, Pace G, Ranieri G, Vicentini C. (2008) May antioxidant therapy improve sperm parameter of men with persistent oligospermia after retrograde embolization for varicocele?. *World J Urol* 26, 97–102.
51. Als Salman ARS., Almashhedy LA. & Hadwan MH., *Biol Trace Elem Res* (2017). <https://doi.org/10.1007/s12011-017-1215-8>