

# Evaluation of Practice of Prescribing and Monitoring Anticoagulants Among Hospitalized Iraqi Patients

Qayssar Joudah Fadheel<sup>1</sup>, Rana Talib Naser<sup>2</sup>, Hayder Abdul-Amir Al-Hindy<sup>3\*</sup>

<sup>1</sup>Department of Pharmacology and Toxicology, College of Pharmacy, University of Babylon, Iraq.

<sup>2</sup>Babylon Health Directorate, [talibrana@gmail.com](mailto:talibrana@gmail.com)

<sup>3</sup>Ph.D., (Medical physiology) College of Pharmacy, University of Babylon

**\*Corresponding Author**

Ass. Prof., Pediatrician, Cardiac Physiology, Department of Pharmacology and Toxicology, University of Pharmacy, Babylon, Iraq.

[phar.hayder.abdul@uobabylon.edu.iq](mailto:phar.hayder.abdul@uobabylon.edu.iq)

Orcid: 0000-0001-6232-8501

Mobile: 00964770802-0200

## Abstract

**Background:** Anticoagulant drugs are lifesaving agents that are prescribed for millions of patients annually. It is essential in the treatment of atrial fibrillation, acute myocardial infarction, as well as deep venous thrombosis. Inaccuracies of their use will lead to life-threatening complications such as bleeding. This study was designed to evaluate and explore the prescribing practice of anticoagulants in the internal medicine ward of Al-Sadr Medical City, Al-Najaf governorate, Iraq.

**Materials and Methods:** The present study is a prospective, survey-based, that was carried out in the wards of internal medicine at Al-Sadr Medical City, during the period from October/2020 to April/2021.

**Results:** 20% of the patients admitted to the ward, were treated with anticoagulants and the rate of their prescribing in respect to all medications administered by the patients was 13%, enoxaparin is being the most commonly prescribed anticoagulant (51%), then unfractionated heparin (40%) and least one was the oral anticoagulant warfarin (9%). Most cases used the anticoagulant drug as monotherapy (92%), and only 8% used combined anticoagulants. The dose range of enoxaparin was about (4000 IU - 6000 IU) and the most frequent dose of enoxaparin reached was (10000 IU), the dose of heparin was about (5000 IU) and for warfarin was 5mg per day. The monitoring of the anticoagulant agents' activity was found to be very rare (8%).

**Conclusion:** The most prescribed anticoagulant was the low molecular weight heparin, and the monitoring of anticoagulants was poor.

**Keywords:** heparin, enoxaparin, warfarin, monitoring, activity, anticoagulants

## Introduction

Throughout healthy homeostasis, the body keeps a steady balance between thrombosis and thrombolysis. This balance is sustained by a multifaceted collaboration between vascular endothelium, platelets, fibrinolytic system, and normal coagulation pathways [1-3]. The normal coagulation cascade (fig-1) encompasses an interface between the tissue factor (extrinsic

pathway) and the contact activation mechanism (intrinsic pathway). These dual pathways apparently independent result to the transformation of "factor X" to active "factor Xa" that is the beginning of the common coagulation pathway. The latter pathway transforms prothrombin to active "thrombin", which is then catalyzes fibrin formation and constantly stabilize the aggregated platelets to form a final stable clot.




Fig 1: Normal coagulation cascade (extrinsic and intrinsic pathways) (1)

[4], and it has been predicted that over 65,000 subjects in U.S. EDs are treated for "warfarin-related hemorrhage. Owing to this bleeding, besides the "narrow therapeutic index" of warfarin and the requirement for repeated monitoring, it is essential to obtain more safe anticoagulant that requires no specific monitoring. Thus, many novel anticoagulant agents created, including direct thrombin inhibitors (dabigatran), factor Xa inhibitors (rivaroxaban, apixaban) and developed to target diverse steps of the coagulation mechanism (fig-1)[2, 5]. Warfarin act by blocking the "vitamin K-epoxide reductase", leading to prevent the synthesis of the "vitamin K-dependent" clotting factors [6, 7]. The recommendations for anticoagulant prescription of "Federal Drug Administration" advocate long-term anticoagulants after any thrombosis or for preventing thrombotic event in high-risk subjects, such as atrial fibrillation, postoperative conditions, and those with artificial cardiac valves [4]. The oral doses of warfarin normally range from 5-10mg/day, adjusted according on the "international normalized ratio (INR)", which is the common global monitoring-index deepened on prothrombin time. The cytochrome P450 system is the main metabolic pathway for [8]. Inhibition or induction of the isozymes that occurred with the metabolism of warfarin can possibly enhance the INR considerably. In addition to that, changes in oral vitamin K intake can develop significant variations in the INR (9). Alternatively, "unfractionated heparin (UFH)" can binds to and enhances the activity of antithrombin-III by causing conformational changes in factor Xa, which inhibit IIa/Xa in a ratio of 1:1 (10). As well, UFH causes some inhibitory effects on factors IXa, XIa, and XIIa [9].

## Materials and methods

### Study design

The present study was a prospective survey-based, carried out in Al-Sadr Medical City, in the internal medicine ward from October/2020 to April/2021.

### Patients

We collect the data for 60 patients taking anticoagulants depending on their records out of 303 total patients who are admitted to the hospital. Thirty-five of them were females and 25 were males. There was a lack of electronic documentation for the data in the hospital. The data were collected from the patient that is why the process takes time to collect these data manually based on the questionnaires, which included (patient personal history, past medical history, diagnosis, type of anticoagulant use, dose regimen, duration, monitoring parameters of prescribed anticoagulant).

### Statistical analysis

The research descriptive statistics mainly used the count and the percentage of anticoagulant use, anticoagulant prescription, and their combination in a pure descriptive statistic. The collected data were summarized and the result was calculated by Microsoft excel.

## Results

### Anticoagulant use and prescribing rate

Out of the total 303 admitted patients, around 20% were receiving anticoagulants (fig-2).




Figure-2: Percent of those were receiving anticoagulants out of the total admitted patients (N-303)

### Anticoagulant prescribing rate

Anticoagulants were prescribed in about a 13% ratio out of the total drugs prescribed to admitted patients (fig-3).




Figure-3: Anticoagulant prescribing rate out of the total drugs prescribed to the admitted patients

In general, females receive anticoagulants more than males, 58% of the anticoagulants were prescribed for female patients, while 42% were prescribed for males (fig-4).




Figure-4: Ratio of female to male receiving anticoagulants among the total included patients

### Type of prescribed anticoagulants

Enoxaparin was the most prescribed anticoagulant (51%), heparin was the second one (40%), and warfarin was the least prescribed one (9%), (fig-5).




Figure-5: Type of prescribed anticoagulants among all admitted patients

### Combined therapy

Most cases were using monotherapy anticoagulants (92%), and only (8%) of the cases used combined anticoagulants (fig-6).




Figure-6: Percent of using combined anticoagulants to monotherapy anticoagulant among admitted patients

### Clinical diagnosis of patients underwent anticoagulant therapy

Out of the total, 35% were on anticoagulants for ischemic heart diseases, 21.6% for heart failure, and 13.3% as part of deep vein thrombosis management or cerebrovascular accidents (fig-7). The remaining were on anticoagulants for treatment of pulmonary embolism (6.7%) or atrial fibrillation and infections (5%).




Figure-7: Clinical diagnosis of the patients that were under anticoagulant therapy

### Therapeutic indications of anticoagulants among the studied patients

Out of the total, 45% (27) were on anticoagulants as primary prophylaxis. 32 (19) were on anticoagulants as treatment and 23% (14) as secondary prophylaxis (fig-8).




Figure-8: Therapeutic indications of anticoagulants among the studied patients

Figure-9 revealed the distribution of the patients according to the usage of the three main anticoagulant therapy based on their therapeutic indications among the included patients. 45% were using anticoagulants as primary prophylaxis, and the remaining 55% were using them either as treatment or secondary prophylaxis (21 and 17), respectively.




Figure-9: Usage of the three main anticoagulant therapy according to their therapeutic indications among the included patients

### Monitoring of the therapy

In about 92% of the patients (fig-10), there was poor monitoring of the activity of anticoagulant agents, this can subsidize further complications.




Figure-10: Percent of monitoring of anticoagulant therapy among included patients

Figure-11 revealed the total daily administered dose of Enoxaparin/IU and Heparin/IU for the admitted patients.




Figure-11: Total daily administered dose of Enoxaparin/IU and Heparin for the admitted patients

### Discussion

To the best of our knowledge, we did not find any previous research concerning anticoagulant prescribing practice in our Iraqi situation. Anticoagulants are life-saving drugs that are prescribed to many patients annually, they are very important in the treatment of deep venous thrombosis [10], acute myocardial infarction [6, 11-13], and atrial fibrillation [14]. Mistakes of their use lead to life-threatening complications like hemorrhage that is why the objective of the current study is to estimate how anticoagulants are practically prescribed, to which patients are usually prescribed, which dose is used, and how it is monitored [9]. We found that about 20% of the patients admitted to the medical ward are treated

with anticoagulants and the prescribing rate of anticoagulant among all medications taken by the patient is 13%, with the low molecular weight heparin (enoxaparin) is being the most prescribed anticoagulant (51%), followed by heparin (40%) and to a lesser extent the oral anticoagulant warfarin (9%). These findings explain the fact which states that low molecular weight heparins are usually preferred because they have a and are effective [15]. The standard prophylactic regimen does not require anticoagulant monitoring [16, 17]. The low molecular weight heparins duration of action is longer than that of heparin (unfractionated) and subcutaneous administration once-daily is possible for some indications, rendering them convenient to use [18]. While warfarin less commonly used has many restrictions due to dosing is variable and its slow onset of action, because its activity is affected by dietary intake of vitamin K and the requirement for frequent and specific monitoring and inherent variability in response with time, the ability for changing the INR potential for causing mistakes in laboratory test values [19].

Most cases use anticoagulants as monotherapy (92%), and only 8% of the cases use a combination. The most frequent dose of enoxaparin reached (10000 IU) and the dose range was about (4000-6000IU). While for heparin the dose was about 5000 IU and for warfarin was 5mg. Anticoagulants are characterized by heterogeneous mechanisms of action and are also different in their effect on coagulation assays [20]. Therefore; an array of assays is required to monitor the activity of anticoagulant drugs which we obtained to be very rare (8%). Such an absence of monitoring can lead to further complications. We observed that in some cases, there is a lack of monitoring even for the critical cases, like patients that take warfarin. Meanwhile, monitoring was observed for less critical cases, such as patients receiving enoxaparin.

Studies have shown that when initiating therapy with warfarin, physicians should leave loading dosages that may increase the INR very high [21]. As a substitute, warfarin must be started with initial dose of 5mg, and in very elderly started with 2-4mg dose, not increase in the initial 24hours, unless in sporadic cases who will continually need quiet small doses (0.5-2.0mg) per day [22]. The adjustment of warfarin dose at a steady-state deepened on the clinical factors and calculated INR measures. The dosage does not require adjustment for a specific INR that is marginally out of range. As well, modifications should adjust the weekly entire dose by 5-20% [23]. The INR requires a frequent monitoring (e. g., 2-4 times/weekly) directly after beginning of warfarin; the intervals between INR measures subsequently, can be extended step by step (reaching to 4-6 weeks) in cases sustained stable INR readings. cases who have a raised INR tests

will necessitate more frequent analysis as well as vitamin K1 supplement [8].

A prior study in Paris revealed that anticoagulant, low molecular weight heparin was used routinely [24]. This study obtained that low molecular weight heparin was prescribed incorrectly to 13% of the patients and in only 26.8% of the patients who are given low molecular weight heparin were fully compliant with all criteria of proper therapeutic use (treatment duration, indications, patient's age, monitoring and renal creatinine clearance).

The findings of the present study were comparable with the outcomes of the reports of the "Public Health Insurance System" in France, which revealed an over-all rate of incorrect anticoagulants prescribing of 47.4% at 1999 [24].

## Conclusions

In the internal medicine ward, the anticoagulant prescription is 13% of the total drugs prescribed to patients, and 20% of the patients admitted were receiving anticoagulants. In general, females receive anticoagulants more than males, 58% of the anticoagulants are prescribed for female patients, while 42% are prescribed for male. Enoxaparin is the most prescribed one (51%), heparin is the second one (40%), and warfarin is the least prescribed one (9%). Most cases use anticoagulants as monotherapy (92%), and only (8%) of the cases use a combination. There is a lack of monitoring of the activity of anticoagulant agents and this can lead to further complications.

## References

1. Abed DA, Jasim Raad, Hayder Abdul-Amir Al-Hindy, Ammar Waheeb Obaid Cystatin-C in patients with acute coronary syndrome: Correlation with ventricular dysfunction, and affected coronary vessels. *Journal of Contemporary Medical Sciences*. 2020;6(1):26-31. <https://doi.org/10.22317/jcms.v6i1.729>. <http://www.jocms.org/index.php/jcms/article/view/729>
2. Amir Al-Mumin HA-AMA-H, Mazin JM. . Combined Assessments of Multi-panel Biomarkers for Diagnostic Performance in Coronary Artery Disease: Case-Control Analysis. *Sys Rev Pharm*. 2020;11(6):7. <https://doi.org/10.31838/ijpr/2020.SP1.193>.
3. Raghdan Z. AKS, Dleikh F., Al-Hindy H. . Is There any Association Between Highly Sensitive C-Reactive Protein and Dental-Status in Ischemic Heart Diseases? A Comparative Stud. *Biochemical and Cellular Archives*. 2020;20(2):6069-75.
4. Flora GD, Nayak MK. A brief review of cardiovascular diseases, associated risk factors and current treatment regimes. *Current pharmaceutical design*. 2019;25(38):4063-84.
5. Hayder Abdul-Amir Maki Al-hindi MJM, Thekra Abid Jaber Al-kashwan, Ahmed Sudan, Saja Ahmed Abdul-Razzaq. On Admission Levels of High Sensitive C- Reactive Protein as A Biomarker in Acute Myocardial Infarction: A Case-Control Study. *Indian Journal of Public Health Research & Development* 2019;10(4):5. <https://doi.org/10.5958/0976-5506.2019.00924.0>.



6. Hajir Karim Abdul-Hussein FSD, Ameera Jasim Al-Aaraji, Hayder Abdul-Amir Makki Al-Hindy, Mazin Jaafar Mousa. Biochemical causal-effect of circulatory uric acid, and HSCRP and their diagnostic correlation in admitted patients with ischemic heart diseases. *Journal of Cardiovascular Disease Research* 2020;11(2):25-31.
7. Thriveni R, Rukhsar I, Ramesh DV, et al. Development and clinical evaluation of transmucosal mucoadhesive patch of lornoxicam for the odontogenic pain management: A preliminary study. *Journal of Natural Science, Biology and Medicine*. 2020;11(1):12-6. [https://doi.org/10.4103/jnsbm.JNSBM\\_224\\_18](https://doi.org/10.4103/jnsbm.JNSBM_224_18).
8. Gage BF, Fihn SD, White RH. Management and dosing of warfarin therapy. *The American Journal of Medicine*. 2000;109(6):481-8. [https://doi.org/10.1016/S0002-9343\(00\)00545-3](https://doi.org/10.1016/S0002-9343(00)00545-3)  
[https://doi.org/10.1016/S0002-9343\(00\)00545-3](https://doi.org/10.1016/S0002-9343(00)00545-3)
9. Palareti G, Antonucci E, Legnani C, et al. Bleeding and thrombotic complications during treatment with direct oral anticoagulants or vitamin K antagonists in venous thromboembolic patients included in the prospective, observational START2-register. 2020;10(11):e040449. <https://doi.org/10.1136/bmjopen-2020-040449>  
<https://bmjopen.bmj.com/content/bmjopen/10/11/e040449.full.pdf>
10. Kearon C, de Wit K, Parpia S, et al. Diagnosis of deep vein thrombosis with D-dimer adjusted to clinical probability: prospective diagnostic management study. 2022;376:e067378. <https://doi.org/10.1136/bmj-2021-067378>  
<https://www.bmj.com/content/bmj/376/bmj-2021-067378.full.pdf>
11. Amir Al-Mumin HA-AMA-H. Hyperuricemia has a Deleterious Role in Patients with Acute Coronary Syndrome Presented with Poor Oral Hygiene. *International Journal of Pharmaceutical Research*. 2020;Jan-Jun(1):7. DOI: <https://doi.org/10.31838/ijpr/2020.SP1.193>.
12. Asseel K. Shaker RA-S, Raad Jasim, Hayder Abdul-Amir Makki Al-Hindy. Biochemical Significance of Cystatin-C and High Sensitive CRP in Patients with Acute Coronary Syndrome; any Clinical Correlation with Diagnosis and Ejection Fraction. *Sys Rev Pharm*. 2020;11(3):8.
13. Hayder Abdul-Amir Maki Al-Hindi MJM, Thekra Abid Jaber Al-Kashwan, Ahmed Sudan, Saja Ahmed Abdul-Razzaq. Correlation of on Admission Levels of Serum Uric Acid with Acute Myocardial Infarction: Case : Control Study. *Journal of Global Pharma Technology*. 2019;11(7):6.
14. Barra S, Providência R. Anticoagulation in atrial fibrillation. 2021;107(5):419-27. <https://doi.org/10.1136/heartjnl-2020-316728>  
<https://heart.bmj.com/content/heartjnl/107/5/419.full.pdf>
15. Ahmed I, Majeed A, Powell R. Heparin induced thrombocytopenia: diagnosis and management update. *Postgrad Med J*. 2007;83(983):575-82. <https://doi.org/10.1136/pgmj.2007.059188>  
<https://pubmed.ncbi.nlm.nih.gov/17823223>  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2600013/>
16. Garcia DA, Baglin TP, Weitz JI, et al. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(2 Suppl):e24S-e43S. <https://doi.org/10.1378/chest.11-2291>.
17. Tran HA, Chunilal SD, Harper PL, et al. An update of consensus guidelines for warfarin reversal. *The Medical journal of Australia*. 2013;198(4):198-9. <https://doi.org/10.5694/mja12.10614>.
18. Solari F VM. Low Molecular Weight Heparin (LMWH) In: StatPearls [Internet]. Treasure Island (FL): Available from: <https://www.ncbi.nlm.nih.gov/books/NBK525957/>. 2022.
19. Weitz JI, Linkins LA. Beyond heparin and warfarin: the new generation of anticoagulants. *Expert opinion on investigational drugs*. 2007;16(3):271-82. <https://doi.org/10.1517/13543784.16.3.271>.
20. Harter K, Levine M, Henderson SO. Anticoagulation drug therapy: a review. *West J Emerg Med*. 2015;16(1):11-7. <https://doi.org/10.5811/westjem.2014.12.22933>  
<https://pubmed.ncbi.nlm.nih.gov/25671002>  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4307693/>
21. Kuruvilla M, Gurk-Turner C. A review of warfarin dosing and monitoring. *Proc (Bayl Univ Med Cent)*. 2001;14(3):305-6. <https://doi.org/10.1080/08998280.2001.11927781>  
<https://pubmed.ncbi.nlm.nih.gov/16369639>  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1305837/>
22. Mansur AdP, Takada JY, Avakian SD, et al. Warfarin doses for anticoagulation therapy in elderly patients with chronic atrial fibrillation. *Clinics (Sao Paulo)*. 2012;67(6):543-6. [https://doi.org/10.6061/clinics/2012\(06\)01](https://doi.org/10.6061/clinics/2012(06)01)  
<https://pubmed.ncbi.nlm.nih.gov/22760890>  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3370303/>
23. Shikdar S VR, Bhattacharya PT. . International Normalized Ratio (INR). Treasure Island (FL), Available from: <https://www.ncbi.nlm.nih.gov/books/NBK507707/>. StatPearls [Internet]. 2021.
24. Tilleul P, Tredan G, Austruy G, et al. Prophylactic low-molecular-weight heparin: Prescription practice in an intensive care unit. *Journal of Critical Care*. 2006;21(2):173-8. <https://doi.org/10.1016/j.jcrc.2005.12.013>  
<https://www.sciencedirect.com/science/article/pii/S0883944106000384>